

# Superconvergence for control-volume mixed finite element methods on rectangular grids

Thomas F. Russell <sup>\*</sup>    Mary F. Wheeler <sup>†</sup>    Ivan Yotov <sup>‡</sup>

## Abstract

We consider control-volume mixed finite element methods for the approximation of second-order elliptic problems on rectangular grids. These methods associate control volumes (covolumes) with the vector variable as well as the scalar, obtaining local algebraic representation of the vector equation (e.g., Darcy's law) as well as the scalar equation (e.g., conservation of mass). We establish  $O(h^2)$  superconvergence for both the scalar variable in a discrete  $L^2$ -norm and the vector variable in a discrete  $H(\text{div})$ -norm. The analysis exploits a relationship between control-volume mixed finite element methods and the lowest order Raviart-Thomas mixed finite element methods.

## 1 Introduction

We consider the second-order elliptic problem in a domain  $\Omega \subset \mathbb{R}^d$ ,  $d = 2$  or  $3$ , written as a first-order system

$$\mathbf{u} = -K\nabla p \text{ in } \Omega, \tag{1.1}$$

$$\nabla \cdot \mathbf{u} = f \text{ in } \Omega, \tag{1.2}$$

$$\mathbf{u} \cdot \mathbf{n} = 0 \text{ on } \partial\Omega. \tag{1.3}$$

The above equations model single-phase flow in porous media, where  $p$  is the fluid pressure, the vector  $\mathbf{u}$  is the Darcy velocity,  $K$  is a symmetric uniformly positive

---

<sup>\*</sup>Department of Mathematics, University of Colorado at Denver, Denver, CO 80217, partially supported by NSF grants DMS 0084438 and DMS 0222300; Division of Mathematical Sciences, National Science Foundation, Arlington, VA 22230, U.S.A.; [trussell@nsf.gov](mailto:trussell@nsf.gov)

<sup>†</sup>Institute for Computational Engineering and Sciences (ICES), Department of Aerospace Engineering & Engineering Mechanics, and Department of Petroleum and Geosystems Engineering, The University of Texas at Austin, Austin, TX 78712, U.S.A.; [mfw@ices.utexas.edu](mailto:mfw@ices.utexas.edu); partially supported by the DOE grant DE-FGO2-04ER25617 and the NSF grants EIA 0121523 and DMS 0411413.

<sup>‡</sup>Department of Mathematics, University of Pittsburgh, Pittsburgh, PA 15260, U.S.A.; [yotov@math.pitt.edu](mailto:yotov@math.pitt.edu); partially supported by the DOE grant DE-FG02-04ER25618 and the NSF grants DMS 0107389 and DMS 0411694.

definite and bounded diagonal tensor with Lipschitz components, representing the rock permeability divided by the fluid viscosity,  $\mathbf{n}$  is the outward unit normal to  $\partial\Omega$ , and  $f$  is the source term satisfying the compatibility condition

$$\int_{\Omega} f \, dx = 0.$$

The choice of homogeneous Neumann boundary condition corresponds to an impermeable boundary, which is the typical physical situation.

In this paper we consider discretizations for (1.1)–(1.3) based on control-volume mixed finite element methods (CVMFEM) and establish  $O(h^2)$  superconvergence for the pressure and velocity in a discrete  $L^2$ -norm and  $H(\text{div})$ -norm, respectively. Most of the arguments can be extended to Dirichlet boundary conditions. However, some loss of superconvergence occurs on the boundary in that case. Global  $O(h)$  convergence has been shown by Chou *et al.* [9, 10]; here we obtain the  $O(h^2)$  rate suggested by various numerical results (e.g., [8, 19, 24, 22]). Superconvergence is proved by  $O(h^2)$  estimates of the differences between the scalar and vector discrete solutions and appropriate projections of the exact solutions.

CVMFEM, first introduced in [8], can be viewed as a type of mixed covolume methods [9, 10, 11]. CVMFEM are closely related to the Raviart-Thomas mixed finite element methods (MFEM) [26, 7, 27], cell-centered finite difference (CCFD) methods [28, 29, 4], mimetic finite difference (MFD) methods [5, 21, 6], and multi-point flux approximation (MPFA) methods [1, 17]. Some of these relationships are explored in detail in [22].

Like MFEM, CVMFEM are designed to provide simultaneous (accurate) approximation of pressure and velocity, and local mass conservation,  $\int_Q \nabla \cdot \mathbf{u}_h = \int_Q f$  on each finite element  $Q$ , where  $\mathbf{u}_h$  is the computed velocity. These properties can be difficult to obtain when  $K$  is heterogeneous (in particular, discontinuous) and/or anisotropic, especially when it incorporates irregular geological features. The methods listed above seek to accomplish this for flow in porous media, among other applications.

Unlike MFEM, CVMFEM have vector control volumes (covolumes) that give rise to a local discrete Darcy law analogous to (1.1). An engineer measuring the permeability of a core sample will typically impose a pressure at each end and observe the flux through the core. The discrete CVMFEM control volume that corresponds to the discrete flux unknown through a face, consisting of the two adjacent halves of the elements on either side of the face (see Figure 1), plays the role of this core, with the element pressures representing the imposed pressures at the ends. The vector test function associated with the control volume is essentially a piecewise-constant vector field, similar to a unit vector in the control volume and a zero vector outside it. The algebraic equation produced by this test function is the local discrete Darcy law. Thus, CVMFEM represent both physical principles in (1.1)–(1.3) locally.

In MFEM, the test vector belongs to the vector trial space, and therefore has continuous normal component. Because the test and trial spaces are the same, the

mass matrix is symmetric and positive definite (SPD). In CVMFEM, the normal component of the test vector is discontinuous at the ends of the control volume, and can also be discontinuous at the element face for general distorted grids. If  $K$  is element-wise constant and the elements are affine (parallelograms in 2D), the mass matrix is SPD, despite the distinct test and trial spaces; in general, it is not symmetric, but symmetry can be restored by appropriate numerical integration [19].

On a uniform grid with homogeneous  $K$ , the lowest-order Raviart-Thomas MFEM, denoted  $\text{RT}_0$ , yields a tridiagonal mass matrix with weights  $1/6$ ,  $2/3$ ,  $1/6$ , and the basic CCFD results in a diagonal mass matrix. As will be seen below, CVMFEM leads to weights  $1/8$ ,  $3/4$ ,  $1/8$ . These are all of the form  $c$ ,  $1 - 2c$ ,  $c$ , where  $c = 0$  (CCFD),  $1/6$  (MFEM), or  $1/8$  (CVMFEM). In [19], some heuristic reasons to favor  $c = 1/8$  are presented: on a uniform grid, the second-order truncation error term is half that of  $c = 0$  and  $c = 1/6$ ; on a nonuniform grid, only  $c = 1/8$  matches one-sided compact finite differences, avoiding any first-order local truncation error; in terms of Fourier modes, the ratio of the discrete eigenvalue to the continuous eigenvalue is generally closer to 1 for  $c = 1/8$ . Numerical results in [22] for homogeneous  $K$  show second-order convergence for both MFEM and CVMFEM; on orthogonal grids, the flux error for CVMFEM improves on that of MFEM by a factor of approximately 2.6; on the distorted grids used, CVMFEM is worse by a factor of about 1.3.

The rest of the paper is organized as follows. In the next section we recall the Raviart-Thomas MFEM for (1.1)–(1.3). Section 3 describes the CVMFEM and its relation to the Raviart-Thomas MFEM. Superconvergence for the velocity is established in Section 4. Section 5 is devoted to superconvergence for the pressure.

## 2 Mixed finite element methods

We will make use of the following standard notation. For a subdomain  $G \subset \mathbb{R}^d$ , the  $L^2(G)$  inner product (or duality pairing) for scalar and vector valued functions is denoted by  $(\cdot, \cdot)_G$ . We denote the norm in the Sobolev space  $W_p^k(G)$ ,  $k \in \mathbb{R}$ ,  $1 \leq p \leq \infty$  [2] by  $\|\cdot\|_{k,p,G}$ . Let  $\|\cdot\|_{k,G}$  be the norm of the Hilbert space  $H^k(G) = W_2^k(G)$ . We omit  $G$  in the subscript if  $G = \Omega$ . For a section of a subdomain boundary  $S \subset \mathbb{R}^{d-1}$  we write  $(\cdot, \cdot)_S$  and  $\|\cdot\|_{0,S}$  for the  $L^2(S)$  inner product (or duality pairing) and norm, respectively.

The mixed variational formulation, which is the basis for the MFEM is: find  $\mathbf{u} \in \mathbf{V}$  and  $p \in W$  such that

$$(K^{-1}\mathbf{u}, \mathbf{v}) = (p, \nabla \cdot \mathbf{v}), \quad \mathbf{v} \in \mathbf{V}, \quad (2.1)$$

$$(\nabla \cdot \mathbf{u}, w) = (f, w), \quad w \in W, \quad (2.2)$$

where

$$\mathbf{V} = \{\mathbf{v} \in H(\text{div}; \Omega) : \mathbf{v} \cdot \mathbf{n} = 0 \text{ on } \partial\Omega\}, \quad W = L_0^2(\Omega) = \left\{ w \in L^2(\Omega) : \int_{\Omega} w \, dx = 0 \right\},$$

and

$$H(\operatorname{div}; \Omega) = \{\mathbf{v} : \mathbf{v} \in (L^2(\Omega))^2, \nabla \cdot \mathbf{v} \in L^2(\Omega)\}$$

with a norm

$$\|\mathbf{v}\|_{\mathbf{V}} = (\|\mathbf{v}\|^2 + \|\nabla \cdot \mathbf{v}\|^2)^{1/2}.$$

We assume that  $\Omega$  can be exactly covered by a rectangular-type finite element partition  $\mathcal{T}_h$ . Let  $\mathbf{V}_h \times W_h \subset \mathbf{V} \times W$  be the lowest order Raviart-Thomas (RT<sub>0</sub>) mixed finite element spaces on  $\mathcal{T}_h$  [26]. More precisely, for all  $Q \in \mathcal{T}_h$ ,

$$\mathbf{V}_h(Q) = \{\mathbf{v} = (a_1 + b_1x, a_2 + b_2y, a_3 + b_3z)^T \text{ on } Q\}, \quad W_h(Q) = \{w = \text{constant on } Q\},$$

$$\mathbf{V}_h = \{\mathbf{v} \in \mathbf{V} : \mathbf{v}|_Q \in \mathbf{V}_h(Q) \forall Q \in \mathcal{T}_h\}, \quad W_h = \{w \in W : w|_Q \in W_h(Q) \forall Q \in \mathcal{T}_h\},$$

where the third component of  $\mathbf{v}$  should be removed if  $d = 2$ . The degrees of freedom of  $\mathbf{V}_h$  are the constant normal components on the sides. If these are continuous, then  $\mathbf{v} \in H(\operatorname{div}; \Omega)$ . Key properties of the RT<sub>0</sub> spaces are

$$\nabla \cdot \mathbf{V}_h = W_h \tag{2.3}$$

and the existence of an interpolation operator  $\Pi : (H^1(\Omega))^d \rightarrow \mathbf{V}_h$  (see [26, 7]) such that for  $\mathbf{q} \in (H^1(\Omega))^2$

$$(\nabla \cdot (\Pi \mathbf{q} - \mathbf{q}), w) = 0, \quad \forall w \in W_h, \tag{2.4}$$

and which satisfies the continuity and approximation properties

$$\|\Pi \mathbf{q}\|_{\mathbf{V}} \leq C \|\mathbf{q}\|_1, \tag{2.5}$$

$$\|\mathbf{q} - \Pi \mathbf{q}\|_0 \leq Ch \|\mathbf{q}\|_1. \tag{2.6}$$

The MFEM for approximating (2.1)–(2.2) is: find  $\tilde{\mathbf{u}}_h \in \mathbf{V}_h, \tilde{p}_h \in W_h$  such that

$$(K^{-1} \tilde{\mathbf{u}}_h, \mathbf{v}) = (\tilde{p}_h, \nabla \cdot \mathbf{v}), \quad \mathbf{v} \in \mathbf{V}_h, \tag{2.7}$$

$$(\nabla \cdot \tilde{\mathbf{u}}_h, w) = (f, w), \quad w \in W_h. \tag{2.8}$$

It has been shown in [26] that (2.7)–(2.8) has a unique solution and

$$\|p - \tilde{p}_h\|_W + \|\mathbf{u} - \tilde{\mathbf{u}}_h\|_{\mathbf{V}} = O(h).$$

A number of authors have studied superconvergence for the above method or the closely related CCFD method [25, 14, 29, 15, 16, 18, 4] and have shown results of the form

$$|||p - \tilde{p}_h|||_W + |||\mathbf{u} - \tilde{\mathbf{u}}_h|||_{\mathbf{V}} = O(h^2),$$

where  $|||\cdot|||_W$  and  $|||\cdot|||_{\mathbf{V}}$  are discrete norms defined in (4.7) and (4.8) below (or some variants of them). The goal of this paper is to obtain similar superconvergence results for the CVMFEM.




Figure 1: Computational grid and control volumes

### 3 Control volume mixed finite element methods

Denote the elements of  $\mathcal{T}_h$  by  $Q_{i,j}$  for  $d = 2$  or by  $Q_{i,j,k}$  for  $d = 3$ , see Figure 1 for  $d = 2$ . For simplicity, in most of the paper we will use the notation and present the arguments for  $d = 2$ . The case  $d = 3$  is a trivial extension.

The center of  $Q_{i,j}$  is denoted by  $c_{i,j}$ . The midpoints of the left and right edges are denoted by  $c_{i-1/2,j}$  and  $c_{i+1/2,j}$ , respectively, with similar notation for the bottom and top edges. With each edge we associate a control volume, where the Darcy's law (1.1) is approximated. In particular, letting  $c_{i+1/2,j} = (x_{i+1/2}, y_j)$ ,  $c_{i,j} = (x_i, y_j)$ , etc., define

$$Q_{i+1/2,j} := (x_i, x_{i+1}) \times (y_{j-1/2}, y_{j+1/2}) \cap \Omega, \quad (3.1)$$

$$Q_{i,j+1/2} := (x_{i-1/2}, x_{i+1/2}) \times (y_i, y_{i+1}) \cap \Omega. \quad (3.2)$$

The control volumes  $Q_{i+1/2,j}$  and  $Q_{i,j+1/2}$  are referred to as  $v_1$ -volumes and  $v_2$ -volumes, respectively. The control volumes that have at least one edge on  $\partial\Omega$  are called border volumes.

Define the velocity test space

$$\mathbf{Y}_h = \{(v_h^1, v_h^2) : \begin{aligned} v_h^1|_{Q_{i+1/2,j}} &= \text{constant} \quad \forall Q_{i+1/2,j}, \quad v_h^1 = 0 \text{ on border } v_1\text{-volumes} \\ v_h^2|_{Q_{i,j+1/2}} &= \text{constant} \quad \forall Q_{i,j+1/2}, \quad v_h^2 = 0 \text{ on border } v_2\text{-volumes} \end{aligned}\}.$$

Thus, for example, the basis function  $\mathbf{y}_{i+1/2,j} \in \mathbf{Y}_h$  associated with  $c_{i+1/2,j}$  is the vector  $(\chi_{i+1/2,j}, 0)$ , i.e.,  $(1, 0)$  on  $Q_{i+1/2,j}$ ,  $(0, 0)$  elsewhere. To see the form of the associated algebraic equation, write (1.1) as  $K^{-1}\mathbf{u} + \nabla p = 0$ , form the inner product with  $\mathbf{y}_{i+1/2,j}$ , and integrate:

$$\int_{x_i}^{x_{i+1}} \int_{y_{j-1/2}}^{y_{j+1/2}} (K^1)^{-1} u^1 dy dx + \int_{y_{j-1/2}}^{y_{j+1/2}} (p(x_{i+1}, y) - p(x_i, y)) dy = 0,$$

where  $\mathbf{u} = (u^1, u^2)$  and  $K = \text{diag}(K^1, K^2)$ . Suppose that  $K$  is element-wise constant on  $Q_{i,j}$  and  $Q_{i+1,j}$ . Taking  $\mathbf{u} = \mathbf{v}_{i-1/2,j}, \mathbf{v}_{i+1/2,j}, \mathbf{v}_{i+3/2,j} \in \mathbf{V}_h$ , the usual RT<sub>0</sub> vector basis functions, we obtain the tridiagonal mass-matrix coefficients

$$1/8 (K_{i,j}^1)^{-1} h_i^x h_j^y, \quad 3/8 (K_{i,j}^1)^{-1} h_i^x h_j^y + 3/8 (K_{i+1,j}^1)^{-1} h_{i+1}^x h_j^y, \quad 1/8 (K_{i+1,j}^1)^{-1} h_{i+1}^x h_j^y,$$

where  $h^x$  and  $h^y$  are the element dimensions. For homogeneous  $K$  and a uniform grid, this reduces to  $1/8, 3/4, 1/8$ , as noted above.

### 3.1 Variational formulation for CVMFEM

Following [9], define the bilinear forms  $a(\cdot, \cdot) : (L^2(\Omega))^d \times (L^2(\Omega))^d \rightarrow \mathbb{R}$ ,  $b(\cdot, \cdot) : \mathbf{Y}_h \times H^1(\Omega) \rightarrow \mathbb{R}$ , and  $c(\cdot, \cdot) : H(\text{div}; \Omega) \times L^2(\Omega) \rightarrow \mathbb{R}$  as follows:

$$\begin{aligned} a(\mathbf{u}, \mathbf{v}) &:= (K^{-1} \mathbf{u}, \mathbf{v}), \\ b(\mathbf{v}, p) &:= \sum_{i,j} \langle p, (v^1, 0)^T \cdot \mathbf{n} \rangle_{\partial Q_{i+1/2,j}} + \sum_{i,j} \langle p, (0, v^2)^T \cdot \mathbf{n} \rangle_{\partial Q_{i,j+1/2}}, \\ c(\mathbf{u}, w) &:= (\nabla \cdot \mathbf{u}, w). \end{aligned}$$

**Lemma 3.1** *If  $(\mathbf{u}, p) \in H(\text{div}; \Omega) \times H^1(\Omega)$  solves (1.1)–(1.3), then  $(\mathbf{u}, p)$  satisfies the variational formulation*

$$a(\mathbf{u}, \mathbf{v}) + b(\mathbf{v}, p) = 0, \quad \mathbf{v} \in \mathbf{Y}_h, \quad (3.3)$$

$$c(\mathbf{u}, w) = (f, w), \quad w \in W_h. \quad (3.4)$$

*Proof:* Equation (1.1) implies, for  $\mathbf{v} \in \mathbf{Y}_h$ ,

$$\begin{aligned} (K^{-1} \mathbf{u}, \mathbf{v}) &= (-\nabla p, \mathbf{v}) = \sum_{i,j} (-\nabla p, (v^1, 0)^T)_{Q_{i+1/2,j}} + \sum_{i,j} (-\nabla p, (0, v^2)^T)_{Q_{i,j+1/2}} \\ &= - \sum_{i,j} \langle p, (v^1, 0)^T \cdot \mathbf{n} \rangle_{\partial Q_{i+1/2,j}} - \sum_{i,j} \langle p, (0, v^2)^T \cdot \mathbf{n} \rangle_{\partial Q_{i,j+1/2}}, \end{aligned}$$

giving (3.3). Equation (3.4) follows trivially from (1.2).  $\square$

The CVMFEM may be formulated as follows: find  $(\mathbf{u}_h, p_h) \in \mathbf{V}_h \times W_h$  such that

$$a(\mathbf{u}_h, \mathbf{v}) + b(\mathbf{v}, p_h) = 0, \quad \mathbf{v} \in \mathbf{Y}_h, \quad (3.5)$$

$$c(\mathbf{u}_h, w) = (f, w), \quad w \in W_h. \quad (3.6)$$

Note that (3.5) is a Petrov-Galerkin FEM, since the test functions differ from the trial functions. We next recall the transfer operator  $\gamma_h : \mathbf{V}_h \rightarrow \mathbf{Y}_h$ , introduced in [9]. Define, for all  $\mathbf{v} \in \mathbf{V}_h$ ,

$$\gamma_h \mathbf{v} = \left( \sum_{i,j} v^1(c_{i+1/2,j}) \chi_{i+1/2,j}, \sum_{i,j} v^2(c_{i,j+1/2}) \chi_{i,j+1/2} \right)$$

It has been shown in [9] that for constants  $\alpha > 0$  and  $C$  independent of  $h$ ,

$$b(\gamma_h \mathbf{v}, w) = -c(\mathbf{v}, w) \quad \forall \mathbf{v} \in \mathbf{V}_h, w \in W_h, \quad (3.7)$$

$$a(\mathbf{v}, \gamma_h \mathbf{v}) \geq \alpha \|\mathbf{v}\|_0^2 \quad \forall \mathbf{v} \in \mathbf{V}_h, \quad (3.8)$$

$$\|\gamma_h \mathbf{v}\|_0 \leq C \|\mathbf{v}\|_0. \quad (3.9)$$

## 4 Velocity superconvergence analysis

In this section we establish superconvergence for the velocity in the CVMFEM. Subtracting (3.5)–(3.6) from (3.3)–(3.4) gives the error equations

$$a(\mathbf{u} - \mathbf{u}_h, \mathbf{v}) + b(\mathbf{v}, p - p_h) = 0, \quad \mathbf{v} \in \mathbf{Y}_h, \quad (4.1)$$

$$c(\mathbf{u} - \mathbf{u}_h, w) = 0, \quad w \in W_h. \quad (4.2)$$

We first note that (4.2) implies

$$0 = c(\mathbf{u} - \mathbf{u}_h, w) = (\nabla \cdot (\mathbf{u} - \mathbf{u}_h), w) = (\nabla \cdot (\Pi \mathbf{u} - \mathbf{u}_h), w), \quad \forall w \in W_h,$$

using (2.4). Therefore, using (2.3),

$$\nabla \cdot (\Pi \mathbf{u} - \mathbf{u}_h) = 0. \quad (4.3)$$

Let  $\hat{p}$  be the  $L^2$ -orthogonal projection of  $p$  onto  $W_h$ , defined by

$$(p - \hat{p}, w) = 0, \quad \forall w \in W_h.$$

Taking  $\mathbf{v} = \gamma_h(\Pi \mathbf{u} - \mathbf{u}_h)$  and  $w = \hat{p} - p_h$  in (4.1)–(4.2) implies

$$\begin{aligned} a(\Pi \mathbf{u} - \mathbf{u}_h, \gamma_h(\Pi \mathbf{u} - \mathbf{u}_h)) \\ = -a(\mathbf{u} - \Pi \mathbf{u}, \gamma_h(\Pi \mathbf{u} - \mathbf{u}_h)) - b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), p - p_h), \end{aligned} \quad (4.4)$$

$$c(\Pi \mathbf{u} - \mathbf{u}_h, \hat{p} - p_h) = 0. \quad (4.5)$$

The second term on the right in (4.4) can be manipulated as follows:

$$\begin{aligned} b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), p - p_h) &= b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), p - \hat{p}) + b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), \hat{p} - p_h) \\ &= b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), p - \hat{p}) - c(\Pi \mathbf{u} - \mathbf{u}_h, \hat{p} - p_h) \\ &= b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), p - \hat{p}), \end{aligned}$$

using (3.7) and (4.5) in the last equality. Therefore (4.4) gives

$$a(\Pi \mathbf{u} - \mathbf{u}_h, \gamma_h(\Pi \mathbf{u} - \mathbf{u}_h)) = -a(\mathbf{u} - \Pi \mathbf{u}, \gamma_h(\Pi \mathbf{u} - \mathbf{u}_h)) - b(\gamma_h(\Pi \mathbf{u} - \mathbf{u}_h), p - \hat{p}) \quad (4.6)$$

Lemma 4.1 below implies that

$$|a(\mathbf{u} - \Pi \mathbf{u}, \gamma_h(\Pi \mathbf{u} - \mathbf{u}_h))| \leq Ch^2 \|\mathbf{u}\|_2 \|\Pi \mathbf{u} - \mathbf{u}_h\|_0.$$

Using (4.3), Lemma 4.2 below gives

$$|b(\gamma_h(\Pi\mathbf{u} - \mathbf{u}_h), p - \hat{p})| \leq Ch^2 \|p\|_3 \|\Pi\mathbf{u} - \mathbf{u}_h\|_0.$$

With the above two bounds and (3.8), (4.6) implies the following superconvergence result.

**Theorem 4.1** *For the CVMFEM approximation  $(\mathbf{u}_h, p_h)$ , there exists a constant  $C$  independent of  $h$  such that*

$$\|\Pi\mathbf{u} - \mathbf{u}_h\|_0 \leq Ch^2 (\|\mathbf{u}\|_2 + \|p\|_3). \quad \square$$

The above result immediately implies superconvergence in the velocity in an  $L^2$  sense along the Gaussian lines. Consider an element  $Q = [a_1, b_1] \times [a_2, b_2]$ . Following [18, 16], for a vector  $\mathbf{q} = (q_1, q_2)$  define

$$\|q_1\|_{1,Q}^2 = (b_2 - a_2) \int_{a_1}^{b_1} \left| q_1 \left( x_1, \frac{a_2 + b_2}{2} \right) \right|^2 dx_1,$$

$$\|q_2\|_{2,Q}^2 = (b_1 - a_1) \int_{a_2}^{b_2} \left| q_2 \left( \frac{a_1 + b_1}{2}, x_2 \right) \right|^2 dx_2,$$

$$\|\mathbf{q}\|^2 = \sum_{i=1}^2 \sum_{Q \in \mathcal{T}_h} \|q_i\|_{i,Q}^2.$$

Note that, for  $\mathbf{q} \in \mathbf{V}_h$ ,  $\|\mathbf{q}\| = \|\mathbf{q}\|_0$ .

**Corollary 4.1** *There exists a constant  $C$  independent of  $h$  such that*

$$\|\mathbf{u} - \mathbf{u}_h\| \leq Ch^2 (\|\mathbf{u}\|_2 + \|p\|_3).$$

*Proof:* It was shown in [16] that

$$\|\mathbf{u} - \Pi\mathbf{u}\| \leq Ch^2 |\mathbf{u}|_2,$$

where  $|\cdot|_2$  denotes the  $H^2$  seminorm. Also, using Theorem 4.1,

$$\|\Pi\mathbf{u} - \mathbf{u}_h\| = \|\Pi\mathbf{u} - \mathbf{u}_h\|_0 \leq Ch^2 (\|\mathbf{u}\|_2 + \|p\|_3).$$

The assertion of the corollary follows from the above two bounds and the triangle inequality.  $\square$

It is also easy to see that  $\nabla \cdot (\mathbf{u} - \mathbf{u}_h)$  is superconvergent at the midpoints of the elements. Define, for a scalar function  $g$ ,

$$\|g\|^2 = \sum_{i,j} |Q_{i,j}| g(c_{i,j})^2. \quad (4.7)$$

Using (4.3) and (2.4),

$$|||\nabla \cdot (\mathbf{u} - \mathbf{u}_h)||| = |||\nabla \cdot (\mathbf{u} - \Pi\mathbf{u})||| = |||\nabla \cdot \mathbf{u} - \widehat{\nabla \cdot \mathbf{u}}||| \leq Ch^2 \|\nabla \cdot \mathbf{u}\|_{2,\infty},$$

where the last inequality follows from Lemma 4.3. Defining

$$|||\mathbf{q}|||_{\mathbf{V}}^2 = |||\mathbf{q}|||^2 + |||\nabla \cdot \mathbf{q}|||^2, \quad (4.8)$$

the above results can be summarized as follows.

**Corollary 4.2** *There exists a constant  $C$  independent of  $h$  such that*

$$|||\mathbf{u} - \mathbf{u}_h|||_{\mathbf{V}} \leq Ch^2(\|\mathbf{u}\|_2 + \|\nabla \cdot \mathbf{u}\|_{2,\infty} + \|p\|_3). \quad \square \quad (4.9)$$

We next proceed with the three lemmas needed in the proof of Theorem 4.1.

**Lemma 4.1** *There exists a constant  $C$  independent of  $h$  such that, for all  $\mathbf{v} \in \mathbf{V}_h$ ,*

$$|a(\mathbf{u} - \Pi\mathbf{u}, \gamma_h \mathbf{v})| \leq Ch^2 \|\mathbf{u}\|_2 \|\mathbf{v}\|_0.$$

*Proof:* We first show that, if  $\mathbf{q} \in (P_1(Q))^2$ , where  $P_k$  is the space of polynomials of degree  $\leq k$ , then

$$\int_Q (\mathbf{q} - \Pi\mathbf{q}) \gamma_h \mathbf{v} \, dx \, dy = 0, \quad \forall \mathbf{v} \in \mathbf{V}_h, \, Q \in \mathcal{T}_h. \quad (4.10)$$

The argument follows the proof of Lemma 3.1 in [16]. Let  $Q = [a, b] \times [c, d]$  and let  $L_1(x)$  and  $\tilde{L}_1(y)$  be the linear Legendre polynomials on  $[a, b]$  and  $[c, d]$ , respectively. It is easy to see that any  $\mathbf{q} \in (P^1(Q))^2$  can be decomposed into

$$\mathbf{q}(x, y) = \bar{\mathbf{q}}(x, y) + (\alpha \tilde{L}_1(y), \beta L_1(x))^T,$$

where  $\bar{\mathbf{q}} \in \mathbf{V}_h(Q)$ . Since  $\bar{\mathbf{q}} - \Pi\bar{\mathbf{q}} = 0$ , it is enough to establish (4.10) for  $\mathbf{q}(x, y) = (\alpha \tilde{L}_1(y), \beta L_1(x))^T$ . It is shown in [16] that in this case  $\Pi\mathbf{q} = 0$ . Therefore

$$\begin{aligned} \int_Q (\mathbf{q} - \Pi\mathbf{q}) \gamma_h \mathbf{v} \, dx \, dy &= \int_Q \mathbf{q} \gamma_h \mathbf{v} \, dx \, dy \\ &= \int_Q (\alpha \tilde{L}_1(y) (\gamma_h \mathbf{v})^1(x, y) + \beta L_1(x) (\gamma_h \mathbf{v})^2(x, y)) \, dx \, dy = 0, \end{aligned}$$

using that for any fixed  $x_0 \in [a, b]$ ,  $(\gamma_h \mathbf{v})^1(x_0, y) \in P_0[c, d]$ , that for any fixed  $y_0 \in [c, d]$ ,  $(\gamma_h \mathbf{v})^2(x, y_0) \in P_0[a, b]$ , and the orthogonality properties of  $L_1(x)$  and  $\tilde{L}_1(y)$ .

We now have

$$\begin{aligned} a(\mathbf{u} - \Pi\mathbf{u}, \gamma_h \mathbf{v}) &= (K^{-1}(\mathbf{u} - \Pi\mathbf{u}), \gamma_h \mathbf{v}) \\ &= \sum_{Q \in \mathcal{T}_h} [K_Q^{-1}(\mathbf{u} - \Pi\mathbf{u}, \gamma_h \mathbf{v})_Q + ((K^{-1} - K_Q^{-1})(\mathbf{u} - \Pi\mathbf{u}), \gamma_h \mathbf{v})_Q], \end{aligned}$$

where  $K_Q^{-1}$  is the value of  $K^{-1}$  at the center of  $Q$ . Therefore, using that  $K^{-1}$  is Lipschitz,

$$|a(\mathbf{u} - \Pi\mathbf{u}, \gamma_h\mathbf{v})| \leq C \sum_{Q \in \mathcal{T}_h} |(\mathbf{u} - \Pi\mathbf{u}, \gamma_h\mathbf{v})_Q| + Ch \|\mathbf{u} - \Pi\mathbf{u}\|_0 \|\gamma_h\mathbf{v}\|_0. \quad (4.11)$$

Using (4.10), an application of the Bramble-Hilbert lemma [12] implies

$$|(\mathbf{u} - \Pi\mathbf{u}, \gamma_h\mathbf{v})_Q| \leq Ch^2 |\mathbf{u}|_{2,Q} \|\gamma_h\mathbf{v}\|_{0,Q},$$

which, combined with (4.11), (2.6), and (3.9) completes the proof.  $\square$

**Lemma 4.2** *There exists a constant  $C$  independent of  $h$  such that, for all  $\mathbf{v} \in \mathbf{V}_h$ ,*

$$|b(\gamma_h\mathbf{v}, p - \hat{p})| \leq Ch^2 \|p\|_3 \|\mathbf{v}\|_{\mathbf{V}}$$

*Proof:* Let  $e_{i+1/2,j} = \partial Q_{i+1/2,j} \cap Q_{i,j}$  and  $e_{i,j+1/2} = \partial Q_{i,j+1/2} \cap Q_{i,j}$ . Note that in the sums in

$$\begin{aligned} & b(\gamma_h\mathbf{v}, p - \hat{p}) \\ &= \sum_{i,j} \langle p - \hat{p}, ((\gamma_h\mathbf{v})^1, 0)^T \cdot \mathbf{n} \rangle_{\partial Q_{i+1/2,j}} + \sum_{i,j} \langle p - \hat{p}, (0, (\gamma_h\mathbf{v})^2)^T \cdot \mathbf{n} \rangle_{\partial Q_{i,j+1/2}}, \end{aligned}$$

every edge  $e_{i+1/2,j}$  and  $e_{i,j+1/2}$  appears twice (from the two neighboring covolumes). Using that  $\frac{\partial v_1}{\partial x}$  and  $\frac{\partial v_2}{\partial y}$  are constants on each element  $Q_{i,j}$ , we have

$$\begin{aligned} & b(\gamma_h\mathbf{v}, p - \hat{p}) \\ &= \sum_{i,j} \left( h_i^x \frac{\partial v_1}{\partial x} \int_{e_{i+1/2,j}} (p - \hat{p}) dy + h_j^y \frac{\partial v_2}{\partial y} \int_{e_{i,j+1/2}} (p - \hat{p}) dx \right) \\ &= \sum_{i,j} \left( \frac{\partial v_1}{\partial x} \left( h_i^x \int_{e_{i+1/2,j}} p dy - \int_{Q_{i,j}} p dx dy \right) \right. \\ & \quad \left. + \frac{\partial v_2}{\partial y} \left( h_j^y \int_{e_{i,j+1/2}} p dx - \int_{Q_{i,j}} p dx dy \right) \right) \\ &= \sum_{i,j} \left( \left( p, \frac{\partial v_1}{\partial x} \right)_{Q_{i,j}, M_x} - \left( p, \frac{\partial v_1}{\partial x} \right)_{Q_{i,j}} + \left( p, \frac{\partial v_2}{\partial y} \right)_{Q_{i,j}, M_y} - \left( p, \frac{\partial v_2}{\partial y} \right)_{Q_{i,j}} \right), \end{aligned} \quad (4.12)$$

where  $(\cdot, \cdot)_{Q, M_x}$  is the quadrature rule on  $Q$  which uses the midpoint rule in  $x$  and exact integration in  $y$ , and  $(\cdot, \cdot)_{Q, M_y}$  uses exact integration in  $x$  and the midpoint rule in  $y$ . Since the midpoint rule is exact for linear polynomials, the Peano Kernel Theorem [13, Theorem 3.7.1] implies

$$\begin{aligned} & \left( p, \frac{\partial v_1}{\partial x} \right)_{Q_{i,j}, M_x} - \left( p, \frac{\partial v_1}{\partial x} \right)_{Q_{i,j}} = \int_{Q_{i,j}} \varphi(x) \frac{\partial^2 p}{\partial x^2}(x, y) \frac{\partial v_1}{\partial x} dx dy \\ &= \int_{Q_{i,j}} \varphi(x) \frac{\partial^2 p}{\partial x^2}(x, y) \nabla \cdot \mathbf{v} dx dy - \int_{Q_{i,j}} \varphi(x) \frac{\partial^2 p}{\partial x^2}(x, y) \frac{\partial v_2}{\partial y} dx dy \equiv T_1 + T_2 \end{aligned} \quad (4.13)$$

where

$$\varphi(x) = \begin{cases} (x - x_{i-1/2})^2/2, & x_{i-1/2} \leq x \leq x_i \\ (x - x_{i+1/2})^2/2, & x_i \leq x \leq x_{i+1/2} \end{cases}$$

For the first term we have

$$|T_1| \leq Ch^2 \|p\|_{2,Q_{i,j}} \|\nabla \cdot \mathbf{v}\|_{0,Q_{i,j}} \quad (4.14)$$

Integrating by parts in  $T_2$  gives

$$\begin{aligned} T_2 &= \int_{Q_{i,j}} \varphi(x) \frac{\partial^3 p}{\partial x^2 \partial y}(x, y) v_2(x, y) dx dy \\ &\quad - \left( \int_{e_{i,j,t}} - \int_{e_{i,j,b}} \right) \varphi(x) \frac{\partial^2 p}{\partial x^2}(x, y) v_2(x, y) dx \equiv T_{2,1} + T_{2,2} \end{aligned} \quad (4.15)$$

where  $e_{i,j,t}$  and  $e_{i,j,b}$  are the top and the bottom edge of  $Q_{i,j}$ , respectively. For  $T_{2,1}$  we have

$$|T_{2,1}| \leq Ch^2 \|p\|_{3,Q_{i,j}} \|\mathbf{v}\|_{0,Q_{i,j}}. \quad (4.16)$$

For  $T_{2,2}$  we notice that  $v_2$  is continuous across horizontal edges and the assumed regularity of  $p(x, y)$  implies that the trace of  $\frac{\partial^2 p}{\partial x^2}$  is well defined. When summing over all elements, each edge integral will appear twice from the expressions for the two neighboring elements, with opposite signs. Therefore

$$\sum_{i,j} T_{2,2} = 0. \quad (4.17)$$

Combining (4.13)–(4.17) implies

$$\sum_{i,j} \left( \left( p, \frac{\partial v_1}{\partial x} \right)_{Q_{i,j}, M_x} - \left( p, \frac{\partial v_1}{\partial x} \right)_{Q_{i,j}} \right) \leq Ch^2 \|p\|_3 \|\mathbf{v}\|_{\mathbf{v}}.$$

The second error term in (4.12) can be bounded in a similar way. Note that for  $d = 3$ , a similar argument goes through with two terms analogous to  $T_2$ .  $\square$

**Lemma 4.3** *For all  $g \in W_\infty^2$  there exists a constant  $C$  independent of  $h$  such that*

$$\| \|g - \hat{g}\| \| \leq Ch^2 \|g\|_{2,\infty}.$$

*Proof:* Let  $Q \in \mathcal{T}_h$ . The Taylor expansion about the midpoint  $(x_0, y_0)$  of  $Q$  gives for any  $(x, y) \in Q$

$$g(x, y) = g(x_0, y_0) + (x - x_0) \frac{\partial g}{\partial x}(x_0, y_0) + (y - y_0) \frac{\partial g}{\partial y}(x_0, y_0) + R(x, y),$$

where  $|R(x, y)| \leq Ch^2 \|g\|_{2, \infty, Q}$ . Integrating the above equation over  $Q$  and using that  $\int_Q g = \int_Q \hat{g}$  gives

$$|Q|(\hat{g}(x_0, y_0) - g(x_0, y_0)) = \int_Q R(x, y) dx dy,$$

which implies

$$|\hat{g}(x_0, y_0) - g(x_0, y_0)| \leq Ch^2 \|g\|_{2, \infty, Q}.$$

The statement of the lemma now follows from the definition (4.7) of  $||| \cdot |||$ .  $\square$

## 5 Pressure superconvergence analysis

In this section we employ a duality argument to derive superconvergence for the pressure at the cell centers. We will make use of the following continuity property of  $\Pi$  [23, 3]: for any  $\varepsilon > 0$ ,

$$\|\Pi \mathbf{q}\|_0 \leq C(\|\mathbf{q}\|_\varepsilon + \|\nabla \cdot \mathbf{q}\|_0). \quad (5.1)$$

Consider the auxiliary problem

$$\begin{aligned} -\nabla \cdot K \nabla \varphi &= \hat{p} - p_h \quad \text{in } \Omega, \\ -K \nabla \varphi \cdot \mathbf{n} &= 0 \quad \text{on } \partial \Omega, \end{aligned} \quad (5.2)$$

which is well posed since  $\int_\Omega \hat{p} = \int_\Omega p_h = 0$ . Elliptic regularity [20] implies that there exists  $\varepsilon > 0$  such that

$$\|\varphi\|_{1+\varepsilon} \leq C\|\hat{p} - p_h\|_0. \quad (5.3)$$

Note that (5.3) holds for L-shaped domains. Let  $\phi = -K \nabla \varphi$ . We have

$$\begin{aligned} \|\hat{p} - p_h\|_0^2 &= (\hat{p} - p_h, \nabla \cdot \phi) = (\hat{p} - p_h, \nabla \cdot \Pi \phi) = c(\Pi \phi, \hat{p} - p_h) \\ &= -b(\gamma_h \Pi \phi, \hat{p} - p_h) = -b(\gamma_h \Pi \phi, \hat{p} - p) - b(\gamma_h \Pi \phi, p - p_h) \\ &= -b(\gamma_h \Pi \phi, \hat{p} - p) + a(\mathbf{u} - \mathbf{u}_h, \gamma_h \Pi \phi), \end{aligned} \quad (5.4)$$

using (4.1) with  $\mathbf{v} = \gamma_h \Pi \phi$ . By Lemma 4.2,

$$|b(\gamma_h \Pi \phi, \hat{p} - p)| \leq Ch^2 \|p\|_3 \|\Pi \phi\|_{\mathbf{v}} \leq Ch^2 \|p\|_3 (\|\phi\|_\varepsilon + \|\nabla \cdot \phi\|_0) \leq Ch^2 \|p\|_3 \|\hat{p} - p_h\|_0,$$

using (5.1), (5.3), and that  $\|\nabla \cdot \Pi \phi\|_0 \leq \|\nabla \cdot \phi\|_0$ , which follows from  $\nabla \cdot \Pi \phi = \widehat{\nabla \cdot \phi}$ . Note that the constant  $C$  above depends on  $\|K\|_{\varepsilon, \infty}$ . For the last term in (5.4) we write

$$\begin{aligned} |a(\mathbf{u} - \mathbf{u}_h, \gamma_h \Pi \phi)| &= |a(\mathbf{u} - \Pi \mathbf{u}, \gamma_h \Pi \phi) + a(\Pi \mathbf{u} - \mathbf{u}_h, \gamma_h \Pi \phi)| \\ &\leq C(h^2 \|\mathbf{u}\|_2 \|\Pi \phi\|_0 + \|\Pi \mathbf{u} - \mathbf{u}_h\|_0 \|\gamma_h \Pi \phi\|_0) \\ &\leq Ch^2 (\|\mathbf{u}\|_2 + \|p\|_3) \|\Pi \phi\|_0 \\ &\leq Ch^2 (\|\mathbf{u}\|_2 + \|p\|_3) \|\hat{p} - p_h\|_0, \end{aligned}$$

using Lemma 4.1, Theorem 4.1, (3.9), (5.1), (5.3), and (5.2). A combination of (5.4) and the above two bounds gives the following pressure superconvergence result.

**Theorem 5.1** *For the CVMFEM approximation  $(\mathbf{u}_h, p_h)$ , there exists a constant  $C$  dependent on  $\|K\|_{\varepsilon, \infty}$ , but independent of  $h$  such that*

$$\|\hat{p} - p_h\|_0 \leq Ch^2(\|\mathbf{u}\|_2 + \|p\|_3). \quad \square$$

It is now easy to obtain superconvergence for the pressure at the midpoints of the elements. Let  $|||w|||_W = |||w|||$ , where  $|||w|||$  is defined in (4.7) and note that  $|||w|||_W = \|w\|_0$  for all  $w \in W_h$ .

**Corollary 5.1** *There exists a constant  $C$  dependent on  $\|K\|_{1, \infty}$ , but independent of  $h$  such that*

$$|||p - p_h|||_W \leq Ch^2(\|\mathbf{u}\|_2 + \|p\|_{2, \infty} + \|p\|_3).$$

*Proof:* The result follows immediately from the triangle inequality

$$|||p - p_h|||_W \leq |||p - \hat{p}|||_W + |||\hat{p} - p_h|||_W,$$

Lemma 4.3, and Theorem 5.1.  $\square$

## References

- [1] I. AAVATSMARK, T. BARKVE, O. BØE, AND T. MANNSETH, *Discretization on unstructured grids for inhomogeneous, anisotropic media. I. Derivation of the methods*, SIAM J. Sci. Comput., 19 (1998), pp. 1700–1716 (electronic).
- [2] R. A. ADAMS, *Sobolev spaces*, Academic Press, New York, 1975.
- [3] T. ARBOGAST, L. C. COWSAR, M. F. WHEELER, AND I. YOTOV, *Mixed finite element methods on non-matching multiblock grids*, SIAM J. Numer. Anal., 37 (2000), pp. 1295–1315.
- [4] T. ARBOGAST, M. F. WHEELER, AND I. YOTOV, *Mixed finite elements for elliptic problems with tensor coefficients as cell-centered finite differences*, SIAM J. Numer. Anal., 34 (1997), pp. 828–852.
- [5] M. BERNDT, K. LIPNIKOV, D. MOULTON, AND M. SHASHKOV, *Convergence of mimetic finite difference discretizations of the diffusion equation*, East-West J. Numer. Math., 9 (2001), pp. 265–284.
- [6] M. BERNDT, K. LIPNIKOV, M. SHASHKOV, M. F. WHEELER, AND I. YOTOV, *Superconvergence of the velocity in mimetic finite difference methods on quadrilaterals*. SIAM J. Numer. Anal., to appear.

- [7] F. BREZZI AND M. FORTIN, *Mixed and hybrid finite element methods*, Springer-Verlag, New York, 1991.
- [8] Z. CAI, J. E. JONES, S. F. MCCORMICK, AND T. F. RUSSELL, *Control-volume mixed finite element methods*, *Comput. Geosci.*, 1 (1997), pp. 289–315.
- [9] S.-H. CHOU AND D. Y. KWAK, *Mixed covolume methods on rectangular grids for elliptic problems*, *SIAM J. Numer. Anal.*, 37 (2000), pp. 758–771 (electronic).
- [10] S.-H. CHOU, D. Y. KWAK, AND K. Y. KIM, *A general framework for constructing and analyzing mixed finite volume methods on quadrilateral grids: the overlapping covolume case*, *SIAM J. Numer. Anal.*, 39 (2001), pp. 1170–1196 (electronic).
- [11] S.-H. CHOU AND P. S. VASSILEVSKI, *A general mixed covolume framework for constructing conservative schemes for elliptic problems*, *Math. Comp.*, 68 (1999), pp. 991–1011.
- [12] P. G. CIARLET, *The finite element method for elliptic problems*, North-Holland, New York, 1978.
- [13] P. J. DAVIS, *Interpolation and Approximation*, Dover Publications, New York, 1975.
- [14] J. DOUGLAS, JR. AND F. A. MILNER, *Interior and superconvergence estimates for mixed methods for second order elliptic problems*, *RAIRO Modél. Math. Anal. Numér.*, 19 (1985), pp. 397–428.
- [15] J. DOUGLAS, JR. AND J. WANG, *Superconvergence of mixed finite element methods on rectangular domains*, *Calcolo*, 26 (1989), pp. 121–133 (1990).
- [16] R. DURÁN, *Superconvergence for rectangular mixed finite elements*, *Numer. Math.*, 58 (1990), pp. 287–298.
- [17] M. G. EDWARDS AND C. F. ROGERS, *Finite volume discretization with imposed flux continuity for the general tensor pressure equation*, *Comput. Geosci.*, 2 (1998), pp. 259–290 (1999).
- [18] R. E. EWING, R. D. LAZAROV, AND J. WANG, *Superconvergence of the velocity along the Gauss lines in mixed finite element methods*, *SIAM J. Numer. Anal.*, 28 (1991), pp. 1015–1029.
- [19] V. A. GARANZHA AND V. N. KONSHIN, *Approximation schemes and discrete well models for the numerical simulation of the 2-d non-darcy fluid flows in porous media*, tech. rep., *Comm. Appl. Math.*, Computer Centre, Russian Academy of Sciences, Moscow, 1999.

- [20] P. GRISVARD, *Elliptic problems in nonsmooth domains*, Pitman, Boston, 1985.
- [21] J. HYMAN, J. MOREL, M. SHASHKOV, AND S. STEINBERG, *Mimetic finite difference methods for diffusion equations*, *Comput. Geosci.*, 6 (2002), pp. 333–352.
- [22] R. A. KLAUSEN AND T. F. RUSSELL, *Relationships among some locally conservative discretization methods which handle discontinuous coefficients*, *Comput. Geosci.*, 8 (2004), pp. 341–377.
- [23] T. P. MATHEW, *Domain decomposition and iterative refinement methods for mixed finite element discretizations of elliptic problems*, PhD thesis, Courant Institute of Mathematical Sciences, New York University, 1989. Tech. Rep. 463.
- [24] R. L. NAFF, T. F. RUSSELL, AND J. D. WILSON, *Shape functions for velocity interpolation in general hexahedral cells*, *Comput. Geosci.*, 6 (2002), pp. 285–314.
- [25] M. NAKATA, A. WEISER, AND M. F. WHEELER, *Some superconvergence results for mixed finite element methods for elliptic problems on rectangular domains*, in *The Mathematics of Finite Elements and Applications V*, J. R. Whiteman, ed., Academic Press, London, 1985, pp. 367–389.
- [26] R. A. RAVIART AND J. M. THOMAS, *A mixed finite element method for 2nd order elliptic problems*, in *Mathematical Aspects of the Finite Element Method*, *Lecture Notes in Mathematics*, vol. 606, Springer-Verlag, New York, 1977, pp. 292–315.
- [27] J. E. ROBERTS AND J.-M. THOMAS, *Mixed and hybrid methods*, in *Handbook of Numerical Analysis*, Vol. II, P. G. Ciarlet and J. Lions, eds., Elsevier Science Publishers B.V., 1991, pp. 523–639.
- [28] T. F. RUSSELL AND M. F. WHEELER, *Finite element and finite difference methods for continuous flows in porous media*, in *The Mathematics of Reservoir Simulation*, R. E. Ewing, ed., SIAM, Philadelphia, 1983, pp. 35–106.
- [29] A. WEISER AND M. F. WHEELER, *On convergence of block-centered finite-differences for elliptic problems*, *SIAM J. Numer. Anal.*, 25 (1988), pp. 351–375.