

**“THE MAZUR PRODUCT OF CONVERGENT SEQUENCES
IS NOT ONTO. 2.”**

FUNCTIONAL ANALYSIS SEMINAR

TUESDAY 11/FEBRUARY/2020, 2:00 PM - 2:50 PM
ROOM 321 THACKERAY

CHRIS LENNARD

Date: February 7, 2020.

Abstract. We present a discussion of a 2009 solution of Mazur's Problem 8 from the Scottish Book by [L/Dan Radelet, JMAA 350, pp. 384-392]. It is well known that the Cesàro mapping C on *the space of convergent sequences of complex numbers* c ,

$$C : x = (x_n)_{n \geq 0} \mapsto \left(\frac{x_0 + x_1 + \cdots + x_n}{n + 1} \right)_{n \geq 0},$$

preserves the limit $L = \lambda(x) := \lim_{n \rightarrow \infty} x_n$, but is **not** onto.

Put differently, C maps c into c , $\lambda(Cx) = \lambda(x)$, and there exists $z = (z_n)_{n \geq 0}$ in c such that $z \neq Cx$, for all $x \in c$. Indeed,

$$z = (1, 0, 1/3, 0, 1/5, 0, 1/7, 0, \dots)$$

is such a sequence.

In the Scottish Book, Stanisław Mazur considered an extension of the Cesàro mapping to a bilinear product \boxtimes on c : for all $x, y \in c$,

$$x \boxtimes y := \left(\frac{x_0 y_n + x_1 y_{n-1} + \cdots + x_n y_0}{n + 1} \right)_{n \geq 0}.$$

It turns out that for all $x, y \in c$, $x \boxtimes y \in c$ and

$$\lambda(x \boxtimes y) = \lambda(x) \lambda(y).$$

Note also that $x \boxtimes \mathbb{I} = Cx$, where $\mathbb{I} := (1, 1, 1, \dots, 1, \dots)$. In the Scottish Book, Problem 8, Professor Mazur asked *whether the mapping $\boxtimes : c \times c \longrightarrow c$ is onto?* The prize for solution, written in the margin, was *5 small beers*.

This question was solved in the *negative* in the 1980's:

[*There exists $z \in c$ such that $z \neq x \boxtimes y$, for all $x, y \in c$*]. It was solved independently by P.P.B. Eggermont and Y.J. Leung, as well as S. Kwapien and A. Pełczyński, and also by V.V. Peller.

Later, L/Dan Radelet (2009) came up with another solution to Mazur's problem 8, by solving a more general question. Note that to solve Mazur's problem in the negative, it is sufficient to showing that $\boxtimes : c \times c_0 \longrightarrow c_0$ is not onto. Here, c_0 is the space of all real sequences that converge to zero. L/Radelet showed that [*There exists $z \in c_0$ such that $z \neq x \boxtimes y$, for all $x \in \mathcal{H}^2$ and for all $y \in \mathcal{H}_0^2$*], where \mathcal{H}^2 is a larger space than c and \mathcal{H}_0^2 is a larger space than c_0 .

We will discuss all this, and the ideas behind our solution (including plenty of illustrative examples), over the span of a few Functional Analysis Seminars.